PATIENT PACKAGE INSERT IN ACCORDANCE WITH THE PHARMACISTS' REGULATIONS (PREPARATIONS) - 1986

The dispensing of this medicine requires a physician's prescription
Read this package insert carefully in its entirety before using this medicine
The format of this leaflet was determined by the Ministry of Health and its content
was checked and approved in February 2010



Augmentin® 250 mg Tablets

Augmentin 250 mg Tablets
Composition:
Active ingredient and quantity:
Each tablet contains:
Amoxicillin (as Trihydrate) 250 mg
Clavulanic acid (as Potassium salt) 125 mg
Inactive ingredients:
Each tablet also contains about 0.55 mg sodium
Core: Colloidal Silica, Sodium Starch Glycolate,
Magnesium Stearate, Microcrystalline Cellulose
Film-Coat: Titanium Dioxide, Hydroxypropyl
Methylcellulose (Methocel E5), Hydroxypropyl
Methylcellulose (Methocel E5), Polyethylene Glycol
4000, Polyethylene Glycol 6000, Silicon Oil
(Dimethicone 500)

Augmentin® 500 mg Tablets

Composition:
Active ingredient and quantity:
Each tablet contains:
Amoxicillin (as Trihydrate) 500 mg
Clavulanic acid (as Potassium salt) 125 mg
Inactive ingredients:
Each tablet also contains about 0.88 mg sodium Core: Colloidal Silica, Sabout 0.88 mg sodium Core: Colloidal Silica, Sodium Starch Glycolate, Magnesium Stearate, Microcrystalline Cellulose Film-Coat: Titanium Dioxide, Hydroxypropyl Methylcellulose (Methocel E5), Hydroxypropyl Methylcellulose (Methocel E15), Polyethylene Glycol 4000, Polyethylene Glycol 6000, Silicon Oil (Dimethicone 500)

* Augmentin® 875 mg Tablets

Composition:
Active ingredient and quantity:
Each tablet contains:
Amoxicillin (as Trihydrate) 875 mg
Clavulanic acid (as Potassium salt) 125 mg
Inactive ingredients:
Each tablet also contains about 1.22 mg sodium
Core: Colloidal Anhydrous Silica, Sodium Starch
Glycolate, Magnesium Stearate, Microcrystalline
Cellulose

Cellulose

Cellulose Film-Coat: Titanium Dioxide, Hydroxypropyl Methylcellulose Scps, Hydroxypropyl Methylcellulose 15cps, Polyethylene Glycol 4000, Polyethylene Glycol 6000, Silicon Oil (Dimethicone 500)

Therapeutic group: Amoxicillin - An antibiotic from the penicillin family. Clavulanic acid - Beta lactamase enzyme inhibitor.

Therapeutic activity:
Antibiotic from the semisynthetic penicillin group intended for the treatment of infections caused by bacteria susceptible to this preparation.

Augmentin® Suspension 125 mg/5 ml

Composition:
Active ingredient and concentration:
Each 5 ml contains:
Amoxicillin (as Trihydrate) 125 mg
Clavulanic acid (as Potassium salt) 31.25 mg

Clavulariic acid (as Potassium sait) 31.25 mg Inactive ingredients: Each 5 ml also contains 12.5 mg aspartame (=phenylalanine 7 mg) Xanthan Gum, Hydroxypropyl Methylcellulose, Aspartame, Succinic Acid, Silica Gel, Colloidal Silica, Raspberry Dry Flavour, Orange Dry Flavour 1, Orange Dry Flavour 2, Golden Syrup Dry Flavour

Augmentin® Suspension 250 mg/5 ml

Composition:
Active ingredient and concentration:
Each 5 ml contains:
Amoxicillin (as Trihydrate) 250 mg
Clavulanic acid (as Potassium salt) 62.50 mg
Inactive ingredients:
Each 5 ml also contains 12.5 mg aspartame (=phenylalanine 7 mg)
Xanthan Gum, Hydroxypropyl Methylcellulose, Aspartame, Succinic
Acid, Silica Gel, Colloidal Silica, Raspberry Dry Flavour, Orange
Dry Flavour 1, Orange Dry Flavour 2, Golden Syrup Dry Flavour

Augmentin® Suspension 400 mg/5 ml

Composition:
Active ingredient and concentration:
Each 5 ml contains:
Amoxicillin (as Trihydrate) 400 mg
Clavulanic acid (as Potassium salt) 57 mg
Inactive ingredients:
Each 5 ml also contains about 1.41 mg sodium
and 16.64 mg aspartame (=phenylalanine 9 mg)
Xanthan Gum, Aspartame, Colloidal Anhydrous Silica, Silicon
Dioxide, Crospovidone, Carmellose Sodium, Magnesium Stearate,
Sodium Benzoate, Strawberry Flavour

When should the preparation not be used?

Do not use this medicine if you have a known sensitivity to any of its ingredients or to other medicines from the penicillin or cephalosporin group.

Do not use if you have suffered in the past from jaundice or liver dysfunction associated with the use of Augmentin or penicillin.

ANTIBIOTICS >ELANA LAHAM *****4906 מבוטח:כחם אילנ<u>ה</u>. TAB AUGMENTIN BID 875 TAB 875mg טבכיות 2 X ביום 14 X 'מים לקחת עם האוכל או אחרי האוכל 34.00: מות :34.00 אכתשלום: 34.00

00100185 :חג'אוי מוחמד 08/09/16↔ Sept 8 2016 מנפק : הנאים נרמין